


## CLAIMS

EPO -DG 1

1. The compound of the general formula (1):

04. 10. 2004



114

5

wherein

W, X and Y are all CH and Z is N; R is halo;

R<sup>1</sup> is aryl, heteroaryl, morpholino, piperidino or pyrrolidino; R<sup>2</sup> is NR<sup>3</sup>R<sup>4</sup>,

R<sup>3</sup> and R<sup>4</sup> are independently H, C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl, C<sub>2-8</sub> alkynyl, aryl,

aryl(C<sub>1-8</sub>)alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl, hetero-

10

aryl(C<sub>1-8</sub>)alkyl, NR<sup>5</sup>R<sup>6</sup>, provided that not both R<sup>3</sup> and R<sup>4</sup> are H or NR<sup>5</sup>R<sup>6</sup>, or

R<sup>3</sup> and R<sup>4</sup> together form a C<sub>3-7</sub> alkylene or C<sub>3-7</sub> alkenylene chain optionally

substituted with one or more C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy groups, or,

together with the nitrogen atom to which they are attached, R<sup>3</sup> and R<sup>4</sup> form a

morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring

15

or a piperazine or piperazine N-(C<sub>1-4</sub>)alkyl (especially N-methyl) ring; and

R<sup>5</sup> and R<sup>6</sup> are independently H, C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl, C<sub>2-8</sub> alkynyl, aryl,

aryl(C<sub>1-8</sub>)alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl or

hetero-aryl(C<sub>1-8</sub>)alkyl;

any of the foregoing alkyl, alkenyl, alkynyl or cycloalkyl groups or moieties (other

20

than for R<sup>6</sup>) being optionally substituted with halogen, cyano, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> al-

kylcarbonyl, C<sub>1-6</sub> alkoxycarbonyl, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylthio, tri(C<sub>1-4</sub>)alkylsilyl,

C<sub>1-6</sub>alkylamino or C<sub>1-6</sub> dialkylamino,

any of the foregoing morpholine, thiomorpholine, piperidine, piperazine and

pyrrolidine rings being optionally substituted with C<sub>1-4</sub> alkyl (especially methyl), and

25

any of the foregoing aryl or heteroaryl groups or moieties being optionally substituted

with one or more substituents selected from halo, hydroxy, mercapto, C<sub>1-6</sub> alkyl,

C<sub>2-6</sub>alkenyl, C<sub>2-6</sub> alkynyl, C<sub>1-6</sub> alkoxy, C<sub>2-6</sub> alkenyloxy, C<sub>2-6</sub> alkynyloxy, halo(C<sub>1-6</sub>)alkyl,

halo(C<sub>1-6</sub>)alkoxy, C<sub>1-6</sub> alkylthio, halo(C<sub>1-6</sub>)alkylthio, hydroxy(C<sub>1-6</sub>)alkyl,

C<sub>1-4</sub>alkoxy(C<sub>1-6</sub>)alkyl, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenoxy, benzyloxy,

30

benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro, -NR<sup>'''</sup>R<sup>'''</sup>,

-NHCOR<sup>'''</sup>, -NHCONR<sup>'''</sup>R<sup>'''</sup>, -CONR<sup>'''</sup>R<sup>'''</sup>, -SO<sub>2</sub>R<sup>'''</sup>, -OSO<sub>2</sub>R<sup>'''</sup>, -COR<sup>'''</sup>, -CR<sup>'''</sup>=NR<sup>'''</sup> or

-N=CR<sup>'''</sup>R<sup>'''</sup>, in which R<sup>'''</sup> and R<sup>'''</sup> are independently hydrogen, C<sub>1-4</sub> alkyl, halo-

(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy, halo(C<sub>1-4</sub>)alkoxy, C<sub>1-4</sub> alkylthio, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl-

(C<sub>1-4</sub>)alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy.

2. A compound according claim 1 wherein

- 5 R<sup>3</sup> is C<sub>1-8</sub> alkyl, halo(C<sub>1-8</sub>)alkyl, hydroxy(C<sub>1-8</sub>)alkyl, C<sub>1-4</sub> alkoxy(C<sub>1-8</sub>)alkyl, C<sub>1-4</sub> alkoxy-halo(C<sub>1-8</sub>)alkyl, tri(C<sub>1-4</sub>)alkylsilyl(C<sub>1-6</sub>)alkyl, C<sub>1-4</sub> alkylcarbonyl(C<sub>1-8</sub>)alkyl, C<sub>1-4</sub> alkyl-carbonylhalo(C<sub>1-8</sub>)alkyl, phenyl(C<sub>1-4</sub>)alkyl, C<sub>2-8</sub> alkenyl, halo(C<sub>2-8</sub>)alkenyl, C<sub>2-8</sub> alkynyl, C<sub>3-8</sub> cycloalkyl optionally substituted with chloro, fluoro or methyl, C<sub>3-8</sub> cycloalkyl-(C<sub>1-4</sub>)alkyl, phenylamino, piperidino or morpholino, the phenyl ring of phenylalkyl or phenylamino being optionally substituted with one, two or three substituents  
10 selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy and halo(C<sub>1-4</sub>)alkoxy; and R<sup>4</sup> is H, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl or amino, or R<sup>3</sup> and R<sup>4</sup> together form a C<sub>3-7</sub> alkylene or C<sub>3-7</sub> alkenylene chain optionally substituted with methyl, or,  
15 together with the nitrogen atom to which they are attached, R<sup>3</sup> and R<sup>4</sup> form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine N-(C<sub>1-4</sub>)alkyl (especially N-methyl) ring, in which the morpholine or piperazine rings are optionally substituted with methyl.

20 3. A compound according to claim 1 or 2 claims wherein

- R<sup>1</sup> is phenyl optionally substituted with from one to five halogen atoms or with from one to three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy or halo(C<sub>1-4</sub>)alkoxy, pyridyl optionally substituted with from one to four halogen atoms or with from one to three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl,  
25 C<sub>1-4</sub>alkoxy or halo(C<sub>1-4</sub>)alkoxy, 2- or 3-thienyl optionally substituted with from one to three halogen atoms or with from one to three substituents selected from halo, C<sub>1-4</sub>alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy or halo(C<sub>1-4</sub>)alkoxy, or piperidino or morpholino both optionally substituted with one or two methyl groups.

30 4. A compound according to claim 3 wherein R<sup>1</sup> is 2,6-difluorophenyl, 2-fluoro-6-chlorophenyl, 2,5,6-trifluorophenyl, 2,4,6-trifluorophenyl, 2,6-difluoro-4-methoxyphenyl or pentafluorophenyl.

5. A compound according to claim 1 wherein W, X and Y are all CH and Z is N;  
35 R is halo, R<sup>1</sup> is aryl, heteroaryl, morpholino, piperidino or pyrrolidino; R<sup>2</sup> is NR<sup>3</sup>R<sup>4</sup>; R<sup>3</sup> and R<sup>4</sup> are independently H, C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl, C<sub>2-8</sub> alkynyl, aryl,

aryl(C<sub>1-8</sub>)alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl, hetero-  
 aryl(C<sub>1-8</sub>)alkyl, NR<sup>5</sup>R<sup>6</sup>, provided that not both R<sup>3</sup> and R<sup>4</sup> are H or NR<sup>5</sup>R<sup>6</sup>, or  
 R<sup>3</sup> and R<sup>4</sup> together form a C<sub>3-7</sub> alkylene or C<sub>3-7</sub> alkenylene chain optionally  
 substituted with one or more C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy groups, or,  
 5 together with the nitrogen atom to which they are attached, R<sup>3</sup> and R<sup>4</sup> form a  
 morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring  
 or a piperazine or piperazine N-(C<sub>1-4</sub>)alkyl (especially N-methyl) ring; and  
 R<sup>5</sup> and R<sup>6</sup> are independently H, C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl, C<sub>2-8</sub> alkynyl, aryl,  
 aryl(C<sub>1-8</sub>)alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl or hetero-  
 10 aryl(C<sub>1-8</sub>)alkyl;  
 any of the foregoing alkyl, alkenyl, alkynyl or cycloalkyl groups or moieties being  
 optionally substituted with halogen, cyano, C<sub>1-6</sub>alkoxy, C<sub>1-6</sub> alkylcarbonyl, C<sub>1-6</sub> alkoxy-  
 carbonyl, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylthio, tri(C<sub>1-4</sub>)alkylsilyl, C<sub>1-6</sub> alkylamino or  
 C<sub>1-6</sub>dialkylamino, any of the foregoing morpholine, thiomorpholine, piperidine,  
 15 piperazine and pyrrolidine rings being optionally substituted with C<sub>1-4</sub> alkyl (especially  
 methyl), and any of the foregoing aryl, heteroaryl, aryloxy or heteroaryl groups being  
 optionally substituted with one or more substituents selected from halo, hydroxy,  
 mercapto, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub>alkynyl, C<sub>1-6</sub> alkoxy, C<sub>2-6</sub> alkenyloxy,  
 C<sub>2-6</sub>alkynyloxy, halo(C<sub>1-6</sub>)alkyl, halo(C<sub>1-6</sub>)alkoxy, C<sub>1-6</sub> alkylthio, halo(C<sub>1-6</sub>)alkylthio,  
 20 hydroxy(C<sub>1-6</sub>)alkyl, C<sub>1-4</sub> alkoxy(C<sub>1-6</sub>)alkyl, C<sub>3-6</sub>cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl,  
 phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro,  
 -NR<sup>'''</sup>R<sup>'''</sup>, -NHCOR<sup>'''</sup>, -NHCONR<sup>'''</sup>R<sup>'''</sup>, -CONR<sup>'''</sup>R<sup>'''</sup>, -SO<sub>2</sub>R<sup>'''</sup>, -OSO<sub>2</sub>R<sup>'''</sup>, -COR<sup>'''</sup>,  
 -CR<sup>'''</sup>=NR<sup>'''</sup> or -N=CR<sup>'''</sup>R<sup>'''</sup>, in which R<sup>'''</sup> and R<sup>'''</sup> are independently hydrogen,  
 C<sub>1-4</sub>alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy, halo(C<sub>1-4</sub>)alkoxy, C<sub>1-4</sub> alkylthio, C<sub>3-6</sub> cycloalkyl,  
 25 C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenyl or benzyl, the phenyl and benzyl groups being  
 optionally substituted with halogen, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy.

6. A compound according to claim 1 wherein W, X and Y are all CH and Z is N;  
 R is halo; R<sup>1</sup> is aryl, heteroaryl, morpholino, piperidino or pyrrolidino; R<sup>2</sup> is NR<sup>3</sup>R<sup>4</sup>,  
 30 R<sup>3</sup> is C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>2-4</sub> alkenyl, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl  
 or phenylamino in which the phenyl ring is optionally substituted with one, two or  
 three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy and  
 halo(C<sub>1-4</sub>)alkoxy; and R<sup>4</sup> is H, C<sub>1-4</sub> alkyl or amino, or  
 R<sup>3</sup> and R<sup>4</sup> together form a C<sub>4-6</sub> alkylene chain optionally substituted with C<sub>1-4</sub> alkyl or  
 35 C<sub>1-4</sub> alkoxy, or  
 together with the nitrogen atom to which they are attached, R<sup>3</sup> and R<sup>4</sup> form a

morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine *N*-(C<sub>1-4</sub>)alkyl (especially *N*-methyl) ring; any of the foregoing alkyl, alkenyl, alkynyl or cycloalkyl groups or moieties being optionally substituted with halogen, cyano, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkoxy-carbonyl, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylthio, tri(C<sub>1-4</sub>)alkylsilyl, C<sub>1-6</sub> alkylamino or C<sub>1-6</sub>dialkylamino, any of the foregoing morpholine, thiomorpholine, piperidine, piperazine and pyrrolidine rings being optionally substituted with C<sub>1-4</sub> alkyl (especially methyl), and any of the foregoing aryl or heteroaryl groups or moieties being optionally substituted with one or more substituents selected from halo, hydroxy, mercapto, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>1-6</sub> alkoxy, C<sub>2-6</sub> alkenyloxy, C<sub>2-6</sub>alkynyloxy, halo(C<sub>1-6</sub>)alkyl, halo(C<sub>1-6</sub>)alkoxy, C<sub>1-6</sub> alkylthio, halo(C<sub>1-6</sub>)alkylthio, hydroxy(C<sub>1-6</sub>)alkyl, C<sub>1-4</sub> alkoxy(C<sub>1-6</sub>)alkyl, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro, -NR<sup>'''</sup>R<sup>'''</sup>, -NHCOR<sup>'''</sup>, -NHCONR<sup>'''</sup>R<sup>'''</sup>, -CONR<sup>'''</sup>R<sup>'''</sup>, -SO<sub>2</sub>R<sup>'''</sup>, -OSO<sub>2</sub>R<sup>'''</sup>, -COR<sup>'''</sup>, -CR<sup>'''</sup>=NR<sup>'''</sup> or -N=CR<sup>'''</sup>R<sup>'''</sup>, in which R<sup>'''</sup> and R<sup>'''</sup> are independently hydrogen, C<sub>1-4</sub>alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy, halo(C<sub>1-4</sub>)alkoxy, C<sub>1-4</sub> alkylthio, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy.

7. A compound according to claim 1 wherein one of W, X and Y are all CH and Z is N; R is halo; R<sup>1</sup> is optionally substituted phenyl; R<sup>2</sup> is NR<sup>3</sup>R<sup>4</sup>, R<sup>3</sup> and R<sup>4</sup> are independently H, C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl, C<sub>2-8</sub> alkynyl, aryl, aryl(C<sub>1-8</sub>)alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl, hetero-aryl(C<sub>1-8</sub>)alkyl, NR<sup>5</sup>R<sup>6</sup>, provided that not both R<sup>3</sup> and R<sup>4</sup> are H or NR<sup>5</sup>R<sup>6</sup>, or R<sup>3</sup> and R<sup>4</sup> together form a C<sub>3-7</sub> alkylene or C<sub>3-7</sub> alkenylene chain optionally substituted with one or more C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkoxy groups, or, together with the nitrogen atom to which they are attached, R<sup>3</sup> and R<sup>4</sup> form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine *N*-(C<sub>1-4</sub>)alkyl (especially *N*-methyl) ring; and R<sup>5</sup> and R<sup>6</sup> are independently H, C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl, C<sub>2-8</sub> alkynyl, aryl, aryl(C<sub>1-8</sub>)alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>3-8</sub> cycloalkyl(C<sub>1-6</sub>)alkyl, heteroaryl or hetero-aryl(C<sub>1-8</sub>)alkyl; any of the foregoing alkyl, alkenyl, alkynyl or cycloalkyl groups or moieties being optionally substituted with halogen, cyano, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkoxycarbonyl, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylthio, tri(C<sub>1-4</sub>)alkylsilyl, C<sub>1-6</sub> alkylamino or C<sub>1-6</sub>dialkylamino; any of the foregoing morpholine, thiomorpholine, piperidine,

piperazine and pyrrolidine rings being optionally substituted with C<sub>1-4</sub> alkyl (especially methyl), and any of the foregoing aryl or heteroaryl groups or moieties, including the phenyl group of R<sup>1</sup>, being optionally substituted with one or more substituents selected from halo, hydroxy, mercapto, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>1-6</sub>alkoxy, C<sub>2-6</sub> alkenyloxy, C<sub>2-6</sub> alkynyloxy, halo(C<sub>1-6</sub>)alkyl, halo(C<sub>1-6</sub>)alkoxy, C<sub>1-6</sub>alkylthio, halo(C<sub>1-6</sub>)alkylthio, hydroxy(C<sub>1-6</sub>)alkyl, C<sub>1-4</sub>alkoxy(C<sub>1-6</sub>)alkyl, C<sub>3-6</sub>cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro, -NR<sup>'''</sup>R<sup>'''</sup>, -NHCOR<sup>'''</sup>, -NHCONR<sup>'''</sup>R<sup>'''</sup>, -CONR<sup>'''</sup>R<sup>'''</sup>, -SO<sub>2</sub>R<sup>'''</sup>, -OSO<sub>2</sub>R<sup>'''</sup>, -COR<sup>'''</sup>, -CR<sup>'''</sup>=NR<sup>'''</sup> or -N=CR<sup>'''</sup>R<sup>'''</sup>, in which R<sup>'''</sup> and R<sup>'''</sup> are independently hydrogen, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub>alkoxy, halo(C<sub>1-4</sub>)alkoxy, C<sub>1-4</sub>alkylthio, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen, C<sub>1-4</sub>alkyl or C<sub>1-4</sub>alkoxy.



8. A compound according to claim 1 wherein W, X and Y are all CH and Z is N; R is halo; R<sup>1</sup> is phenyl optionally substituted with from one to five halogen atoms or with from one to three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub>alkoxy or halo(C<sub>1-4</sub>)alkoxy, pyridyl optionally substituted with from one to four halogen atoms or with from one to three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub>alkoxy or halo(C<sub>1-4</sub>)alkoxy, 2- or 3-thienyl optionally substituted with from one to three halogen atoms or with from one to three substituents selected from halo, C<sub>1-4</sub>alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub>alkoxy or halo(C<sub>1-4</sub>)alkoxy, or piperidino or morpholino both optionally substituted with one or two methyl groups; R<sup>2</sup> is NR<sup>3</sup>R<sup>4</sup>; R<sup>3</sup> is C<sub>1-8</sub> alkyl, halo(C<sub>1-8</sub>)alkyl, hydroxy(C<sub>1-8</sub>)alkyl, C<sub>1-4</sub>alkoxy(C<sub>1-8</sub>)alkyl, C<sub>1-4</sub>alkoxy-halo(C<sub>1-8</sub>)alkyl, tri(C<sub>1-4</sub>)alkylsilyl(C<sub>1-6</sub>)alkyl, C<sub>1-4</sub>alkylcarbonyl(C<sub>1-8</sub>)alkyl, C<sub>1-4</sub>alkyl-carbonylhalo(C<sub>1-8</sub>)alkyl, phenyl<sub>(1-4)</sub>alkyl, C<sub>2-8</sub> alkenyl, halo(C<sub>2-8</sub>)alkenyl, C<sub>2-8</sub> alkynyl, C<sub>3-8</sub> cycloalkyl optionally substituted with chloro, fluoro or methyl, C<sub>3-8</sub> cycloalkyl-(C<sub>1-4</sub>)alkyl, phenylamino, piperidino or morpholino, the phenyl ring of phenylalkyl or phenylamino being optionally substituted with one, two or three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub>alkoxy and halo(C<sub>1-4</sub>)alkoxy; and R<sup>4</sup> is H, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl or amino, or R<sup>3</sup> and R<sup>4</sup> together form a C<sub>3-7</sub> alkylene or C<sub>3-7</sub> alkenylene chain optionally substituted with methyl, or, together with the nitrogen atom to which they are attached, R<sup>3</sup> and R<sup>4</sup> form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring

- 92 -

or a piperazine or piperazine N-(C<sub>1-4</sub>)alkyl (especially N-methyl) ring, in which the morpholine or piperazine rings are optionally substituted with methyl.


9. A compound according to claim 1 wherein one of W, X and Y are all CH and Z is N; R is halo; R<sup>1</sup> is phenyl optionally substituted with from one to five halogen atoms or with from one to three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub>alkoxy or halo(C<sub>1-4</sub>)alkoxy; R<sup>2</sup> is NR<sup>3</sup>R<sup>4</sup>; R<sup>3</sup> is C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>2-6</sub>alkenyl, C<sub>3-6</sub> cycloalkyl, C<sub>3-6</sub> cycloalkyl(C<sub>1-4</sub>)alkyl or phenylamino in which the phenyl ring is optionally substituted with one, two or three substituents selected from halo, C<sub>1-4</sub> alkyl, halo(C<sub>1-4</sub>)alkyl, C<sub>1-4</sub> alkoxy and halo(C<sub>1-4</sub>)alkoxy; and R<sup>4</sup> is H, C<sub>1-4</sub> alkyl or amino, or R<sup>3</sup> and R<sup>4</sup> together form a C<sub>4-6</sub> alkylene chain optionally substituted with methyl, or, together with the nitrogen atom to which they are attached, R<sup>3</sup> and R<sup>4</sup> form a morpholine ring.

10. A process for preparing a compound of the general formula (1) according to claim 1 wherein one of R is chloro or fluoro and R<sup>2</sup> is NR<sup>3</sup>R<sup>4</sup> and W, X, Y, Z, R<sup>1</sup>, R<sup>3</sup> and R<sup>4</sup> are as defined in claim 1, which comprises reacting an amine of the general formula NR<sup>3</sup>R<sup>4</sup> with a compound of the general formula (6) or (13):



wherein W, X, Y, Z and R<sup>1</sup> are as defined in claim 1.

~~11. The intermediate chemicals having the general formulae (5), (6) and (13):~~



~~wherein W, X, Y, Z and R<sup>1</sup> are as defined in claim 1.~~

12. A plant fungicidal composition comprising a fungicidally effective amount of a compound as defined in claim 1 and a suitable carrier or diluent therefor.

1-2005

GB0305248

19.APR.2005 16:12

SYNGENTA AG +41 61 3237010

NR.860 S.3

- 93 -

- ~~13.~~ A method of combating or controlling phytopathogenic fungi which comprises  
~~12.~~ applying to a plant, to a seed of a plant, to the locus of the plant or seed or to soil or  
to any other plant growth medium, a fungicidally effective amount of a compound  
according to claim 1 or a composition according to claim 12.